arXiv:1603.08292v1 [hep-ph] 28 Mar 2016

Inclusive three jet production at the LHC as a new BFKL probe

F. Caporale[1], G. Chachamis[1], B.Murdaca[2], A. Sabio Vera[1]

[1] Instituto de Física Teórica UAM/CSIC, Nicolás Cabrera 15
& Universidad Autónoma de Madrid, E-28049 Madrid, Spain.
[2] Dipartimento di Fisica, Università della Calabria &
Istituto Nazionale di Fisica Nucleare, Gruppo Collegato di Cosenza,
I-87036 Arcavacata di Rende, Cosenza, Italy.

March 29, 2016

1 Introduction

The Balitsky-Fadin-Kuraev-Lipatov (BFKL) formalism [1, 2, 3, 4, 5] is one of the most important resummation programmes in high energy QCD. A typical BFKL observable at the LHC is the azimuthal angle (ϕ) decorrelation of two tagged forward/backward jets widely separated in rapidity, Y, in the so-called Mueller-Navelet jets setup [6]. This multiple emission appears as a fast decrease of $\langle \cos(n\,\phi)\rangle$ as a function of Y [7, 8, 9, 10]. However, these differential distributions suffer from a large influence of collinear regions in phase space, so it was proposed to remove the $n = 0$ dependence by studying the ratios $\mathcal{C}_{m,n} = \langle \cos(m\,\phi)\rangle / \langle \cos(n\,\phi)\rangle$ [11, 12, 13]. In recent studies [14, 15], a BFKL analysis at NLL is able to fit the large Y tail of the Mueller-Navelet $\mathcal{C}_{m,n}$ ratios.

In Ref. [16], we proposed new observables related to final states with two tagged forward jets separated by a large rapidity span, along with a third tagged jet produced in the central region of rapidity, allowing for inclusive radiation in the remaining areas of the detectors.

The two tagged forward jets A and B have transverse momentum $\vec{k}_{A,B}$, azimuthal angle $\theta_{A,B}$ and rapidity $Y_{A,B}$. The central jet is characterized by $\vec{k}_J$, θ_J and y_J and the differential cross section on these variables can be written in the form

$$\begin{aligned}\frac{d^3\sigma^{3-\text{jet}}}{d^2\vec{k}_J dy_J} &= \frac{\bar{\alpha}_s}{\pi k_J^2}\int d^2\vec{p}_A \int d^2\vec{p}_B\,\delta^{(2)}\left(\vec{p}_A+\vec{k}_J-\vec{p}_B\right) \\ &\times \varphi\left(\vec{k}_A,\vec{p}_A,Y_A-y_J\right)\varphi\left(\vec{p}_B,\vec{k}_B,y_J-Y_B\right)\end{aligned} \tag{1}$$

where we assume that $Y_A > y_J > Y_B$ and k_J lies above the experimental resolution scale. φ are BFKL gluon Green functions normalized to $\varphi(\vec{p},\vec{q},0) = \delta^{(2)}(\vec{p}-\vec{q})$ and $\bar{\alpha}_s = \alpha_s N_c/\pi$.



Figure 1: The ratios $\mathcal{R}^{1,1}_{2,2}$ and $\mathcal{R}^{1,2}_{2,2}$ as a function of the rapidity of the central jet y_J.

Now, new distributions can be defined using the projections on the two relative azimuthal angles formed by each of the forward jets with the central jet, $\theta_A - \theta_J - \pi$ and $\theta_J - \theta_B - \pi$:

$$\begin{aligned}
&\int_0^{2\pi} d\theta_A \int_0^{2\pi} d\theta_B \int_0^{2\pi} d\theta_J \cos\left(M\left(\theta_A - \theta_J - \pi\right)\right) \cos\left(N\left(\theta_J - \theta_B - \pi\right)\right) \frac{d^3\sigma^{3-\text{jet}}}{d^2\vec{k}_J dy_J} \\
&= \bar{\alpha}_s \sum_{L=0}^{N} \binom{N}{L} \left(k_J^2\right)^{(L-1/2)} \int_0^{\infty} dp^2 \, \left(p^2\right)^{(N-L/2)} \int_0^{2\pi} d\theta \frac{(-1)^{M+N} \cos\left(M\theta\right) \cos\left((N-L)\theta\right)}{\sqrt{\left(p^2 + k_J^2 + 2\sqrt{p^2 k_J^2}\cos\theta\right)^N}} \\
&\quad \times \phi_M\left(p_A^2, p^2, Y_A - y_J\right) \phi_N\left(p^2 + k_J^2 + 2\sqrt{p^2 k_J^2}\cos\theta, p_B^2, y_J - Y_B\right),
\end{aligned} \tag{2}$$

with ϕ_n defined in Ref. [16]. The experimentally relevant observable is the mean value in the selected events of the two cosines, *i.e.*

$$\begin{aligned}
&\langle \cos\left(M\left(\theta_A - \theta_J - \pi\right)\right) \cos\left(N\left(\theta_J - \theta_B - \pi\right)\right) \rangle \\
&= \frac{\int_0^{2\pi} d\theta_A d\theta_B d\theta_J \cos\left(M\left(\theta_A - \theta_J - \pi\right)\right) \cos\left(N\left(\theta_J - \theta_B - \pi\right)\right) \frac{d^3\sigma^{3-\text{jet}}}{d^2\vec{k}_J dy_J}}{\int_0^{2\pi} d\theta_A d\theta_B d\theta_J \frac{d^3\sigma^{3-\text{jet}}}{d^2\vec{k}_J dy_J}}.
\end{aligned} \tag{3}$$

In order to have optimal perturbative convergence and eliminate collinear contamination, we can remove the contributions from zero conformal spin by defining the ratios:

$$\mathcal{R}^{M,N}_{P,Q} = \frac{\langle \cos\left(M\left(\theta_A - \theta_J - \pi\right)\right) \cos\left(N\left(\theta_J - \theta_B - \pi\right)\right) \rangle}{\langle \cos\left(P\left(\theta_A - \theta_J - \pi\right)\right) \cos\left(Q\left(\theta_J - \theta_B - \pi\right)\right) \rangle}, \tag{4}$$

with M, N, P, $Q = 1, 2$.

2 Results and Outlook

Now, different momenta configurations can be investigated. Here, in Fig. 1, two ratios $\mathcal{R}_{P,Q}^{M,N}$ with $M, N = 1, 2$ are shown while the momenta of the forward jets are fixed to $k_A = 40$ GeV and $k_B = 50$ GeV and their rapidities also fixed to $Y_A = 10$ and $Y_B = 0$. The central jet transverse momentum takes three values, $k_J = 30, 45, 70$ GeV and the rapidity of the central jet y_J is varied in between the forward/backward jet rapidities.

It will be very interesting to see the predictions from fixed order analyses as well as from the BFKL inspired Monte Carlo `BFKLex` [17, 18, 19, 20, 21, 22, 23] for these and other similar observables [24, 25] where the projection on azimuthal angles is used. This type of observables will be crucial to define the region of phenomenological applicability of the BFKL resummation.

Acknowledgements

G.C. acknowledges support from the MICINN, Spain, under contract FPA2013-44773-P. A.S.V. acknowledges support from Spanish Government (MICINN (FPA2010-17747, FPA2012-32828)) and, together with F.C and B.M., to the Spanish MINECO Centro de Excelencia Severo Ochoa Programme (SEV-2012-0249). The work of B.M. was supported in part by the grant RFBR-13-02-90907 and by the European Commission, European Social Fund and Calabria Region, that disclaim any liability for the use that can be done of the information provided here.

References

[1] I. I. Balitsky and L. N. Lipatov, "The Pomeranchuk Singularity in Quantum Chromodynamics," *Sov. J. Nucl. Phys.*, **28**(1978), 822, [Yad. Fiz.28,1597(1978)].

[2] E. A. Kuraev, L. N. Lipatov, and Victor S. Fadin, "The Pomeranchuk Singularity in Nonabelian Gauge Theories," *Sov. Phys. JETP*, **45**(1977), 199, [Zh. Eksp. Teor. Fiz.72,377(1977)].

[3] E. A. Kuraev, L. N. Lipatov, and Victor S. Fadin, "Multi - Reggeon Processes in the Yang-Mills Theory," *Sov. Phys. JETP*, **44**(1976), 443, [Zh. Eksp. Teor. Fiz.71,840(1976)].

[4] Victor S. Fadin and L. N. Lipatov, "BFKL pomeron in the next-to-leading approximation," *Phys. Lett.*, **B429**(1998), 127, `hep-ph/9802290`.

[5] Marcello Ciafaloni and Gianni Camici, "Energy scale(s) and next-to-leading BFKL equation," *Phys. Lett.*, **B430**(1998), 349, `hep-ph/9803389`.

[6] Alfred H. Mueller and H. Navelet, "An Inclusive Minijet Cross-Section and the Bare Pomeron in QCD," *Nucl. Phys.*, **B282**(1987), 727.

[7] Vittorio Del Duca and Carl R. Schmidt, "Dijet production at large rapidity intervals," *Phys. Rev.*, **D49**(1994), 4510, hep-ph/9311290.

[8] W. James Stirling, "Production of jet pairs at large relative rapidity in hadron hadron collisions as a probe of the perturbative pomeron," *Nucl. Phys.*, **B423**(1994), 56, hep-ph/9401266.

[9] Lynne H. Orr and W. James Stirling, "Dijet production at hadron hadron colliders in the BFKL approach," *Phys. Rev.*, **D56**(1997), 5875, hep-ph/9706529.

[10] J. Kwiecinski, Alan D. Martin, L. Motyka, and J. Outhwaite, "Azimuthal decorrelation of forward and backward jets at the Tevatron," *Phys. Lett.*, **B514**(2001), 355, hep-ph/0105039.

[11] Agustin Sabio Vera, "The Effect of NLO conformal spins in azimuthal angle decorrelation of jet pairs," *Nucl. Phys.*, **B746**(2006), 1, hep-ph/0602250.

[12] Agustin Sabio Vera and Florian Schwennsen, "The Azimuthal decorrelation of jets widely separated in rapidity as a test of the BFKL kernel," *Nucl. Phys.*, **B776**(2007), 170, hep-ph/0702158.

[13] M. Angioni, G. Chachamis, J. D. Madrigal, and A. Sabio Vera, "Dijet Production at Large Rapidity Separation in N=4 SYM," *Phys. Rev. Lett.*, **107**(2011), 191601, 1106.6172.

[14] B. Ducloue, L. Szymanowski, and S. Wallon, "Evidence for high-energy resummation effects in Mueller-Navelet jets at the LHC," *Phys. Rev. Lett.*, **112**(2014), 082003, 1309.3229.

[15] Francesco Caporale, Dmitry Yu. Ivanov, Beatrice Murdaca, and Alessandro Papa, "MuellerNavelet jets in next-to-leading order BFKL: theory versus experiment," *Eur. Phys. J.*, **C74**(2014) (10), 3084, [Erratum: Eur. Phys. J.C75,no.11,535(2015)], 1407.8431.

[16] F. Caporale, G. Chachamis, B. Murdaca, and A. Sabio Vera, "BFKL predictions for inclusive three jet production at the LHC," *Phys. Rev. Lett.*, **116**(2016) (1), 012001, 1508.07711.

[17] G. Chachamis, M. Deak, A. Sabio Vera, and P. Stephens, "A Comparative study of small x Monte Carlos with and without QCD coherence effects," *Nucl. Phys.*, **B849**(2011), 28, 1102.1890.

[18] G. Chachamis and A. Sabio Vera, "The Colour Octet Representation of the Non-Forward BFKL Green Function," *Phys. Lett.*, **B709**(2012), 301, 1112.4162.

[19] G. Chachamis and A. Sabio Vera, "The NLO N =4 SUSY BFKL Green function in the adjoint representation," *Phys. Lett.*, **B717**(2012), 458, 1206.3140.

[20] G. Chachamis, A. Sabio Vera, and C. Salas, "Bootstrap and momentum transfer dependence in small x evolution equations," *Phys. Rev.*, **D87**(2013) (1), 016007, 1211.6332.

[21] F. Caporale, G. Chachamis, J. D. Madrigal, B. Murdaca, and A Sabio Vera, "A study of the diffusion pattern in N = 4 SYM at high energies," *Phys. Lett.*, **B724**(2013), 127, 1305.1474.

[22] G. Chachamis and A. Sabio Vera, "A Monte Carlo study of double logarithms in the small x region," (2015), 1511.03548.

[23] G. Chachamis and A. Sabio Vera, "The high-energy radiation pattern from BFKLex with double-log collinear contributions," (2015), 1512.03603.

[24] F. Caporale, Francesco Giovanni Celiberto, G. Chachamis, and A. Sabio Vera, "Multi-Regge kinematics and azimuthal angle observables for inclusive four-jet production," (2015), 1512.03364.

[25] F. Caporale, Francesco Giovanni Celiberto, G. Chachamis, D. Gordo Gomez, and A. Sabio Vera, "BFKL Azimuthal Imprints in Inclusive Three-jet Production at 7 and 13 TeV," (2016), 1603.07785.